

# Genetic Improvements in Agriculture

From Hunter Gatherer to Green Revolution  
and Beyond



Cro magnon skull  
*Homo sapiens*  
~ 30,000 years old


- Life on earth is about four billion years old
- *Homo sapiens* emerged as a species about 300,000 years ago



Cro magnon skull  
*Homo sapiens*  
~ 30,000 years old

•Human activities have caused vast changes in the physical, chemical, geological, atmospheric and biological realm of our planet

# We've removed mountains



The Hobet coal mine in Boone County, West Virginia spreads over 10,000 acres (15.6 square miles)

[NASA images](#) by Robert Simmon, based on [Landsat 5](#) data from the USGS [Global Visualization Viewer](#).



# ...dammed rivers...



Three Gorges Dam, the world's largest hydroelectric power generator. The reservoir just upstream of the dam is more than 2 miles (3 km) across



1987



2006

[+ View movie of the dam's construction](#)

Photo credit: NASA/Goddard Space Flight Center Scientific Visualization Studio United States Geological Survey and [Le Grand Portage](#)

# ....caused extinctions.....

The dodo (*Raphus cucullatus*) was a large flightless bird indigenous to Mauritius. The arrival of humans led to the dodo's extinction by the end of the 17<sup>th</sup> century.





# ...modified other species in extraordinary ways





# ... and modified plant genomes for thousands of years.....





# Now we face our biggest challenges



How do we feed more people without further damaging our planet?

Photo courtesy [Earth Observatory NASA](#)

# What is the role of plant breeding in addressing global challenges?



Photo credits: Xochiquetzal Fonseca/[CIMMYT](#) and [IRRI](#)



# GENETIC IMPROVEMENTS IN AGRICULTURE

## **The Distant Past**

Crop plant domestication and beyond

## **The Recent Past**

Hybrid seed

The (First) Green Revolution

Advances in breeding technologies

## **Now and Into The Future**

Breeding for improved human health

Breeding for drought tolerance

Agricultural innovation in Africa


The Second Green Revolution






# The Distant Past (>10,000 years ago to 1900)

*Homo sapiens* originated  
400,000 – 250,000 years ago  
Major crops were domesticated  
~ 10,000 – 5000 years ago  
The development of human  
civilizations is correlated with  
the development of agriculture



[Karol Schauer](#)


# Plant domestication followed the end of the most recent glacial period



Allaby, R.G., Fuller, D.Q., and Brown, T.A. (2008) The genetic expectations of a protracted model for the origins of domesticated crops. Proc. Natl. Acad. Sci. USA 105: [13982-13986](#), copyright National Academy of Sciences USA

# How did people begin to cultivate plants?


It is thought to have been a gradual change from seeking and following food sources






# How did people begin to cultivate plants?

It is thought to have been a gradual change from seeking and following food sources to semi settled migration




# How did people begin to cultivate plants?

It is thought to have been a gradual change from seeking and following food sources to semi settled migration and finally permanent settlements.



# Plants were domesticated in parallel in several regions


Wheat, barley, pea, lentil  
~ 13,000 years ago



Reprinted by permission from Macmillan Publishers Ltd.: [Nature] Diamond, J. (2002). Evolution, consequences and future of plant and animal domestication. Nature 418: [700-707](#), copyright 2002.





# Genetic modification arose as a consequence of cultivation



Planting seeds from “good” plants increased their representation in subsequent generations



Natural variation within population





# During maize domestication cob size increased

Cobs from  
archeological  
sites in the Valley  
of Tehuacan,  
Mexico



Photo © Robert S. Peabody Museum of Archaeology, Phillips Academy, Andover, Massachusetts. All Rights Reserved.

# The hard casings around many grains were eliminated



Teosinte, the wild relative of maize, has hard coverings over each grain. Humans selected against these during maize domestication.



Photo by [Hugh Iltis](#); Reprinted from Doebley, J.F., Gaut, B.S., and Smith, B.D. (2006). The Molecular Genetics of Crop Domestication. Cell 127: [1309-1321](#), with permission from Elsevier.



# Decrease in branching and increase in seed size were also selected for





Image credit Nicolle Rager Fuller,  
[National Science Foundation](#)

# Seeds that don't break off were selected



**Wild**  
Shattering grain  
“Brittle rachis”  
Advantage –  
maximizes seed  
dispersal




**Domesticated**  
Non-shattering grain  
“Tough rachis”  
Advantage –  
facilitates harvesting




From Konishi, S., Izawa, T., Lin, S.Y., Ebana, K., Fukuta, Y., Sasaki, T., and Yano, M. (2006). An SNP caused loss of seed shattering during rice domestication. *Science* 312: [1392-1396](#). Reprinted with permission from AAAS.

# Many of our crops are products of extensive genomic rearrangements




Polyploid (multi-genome) plants are often bigger and so selected for propagation



From Dubcovsky, J. and Dvorak, J. (2007). Genome Plasticity a Key Factor in the Success of Polyploid Wheat Under Domestication. *Science*. 316: 1862-1866. Reprinted with permission from AAAS. Brassica figure from [Adenosine](#)




# Domestication through genome modification gave us modern crops



From Dubcovsky, J. and Dvorak, J. (2007). Genome Plasticity a Key Factor in the Success of Polyploid Wheat Under Domestication. Science. 316: 1862-1866. Reprinted with permission from AAAS.

# Breeding tree of Indica Rice IR64


One of the most widely grown crops, indica rice IR64 is the product of a complex breeding program that has caused extensive genomic modification, mutation, deletion and rearrangement



# The myth of natural food



The food we eat comes from plants already extensively modified from their original form. Even heritage varieties are extensively genetically modified.



Credit: [Nicolle Rager Fuller](#), National Science Foundation




# The Recent Past – Scientific Plant Breeding



Google™

The twentieth century took us from gas lamps to Google and steamships to space shuttles




# The Recent Past – Scientific Plant Breeding




Google™

And the world  
population  
quadrupled in just  
over 100 years

World Population (billions)



The twentieth  
century took  
us from gas  
lamps to  
Google and  
steamships to  
space shuttles



# The Recent Past – Scientific Plant Breeding



Improvements in plant propagation and breeding were needed to keep up with population growth









Photo credits: [Gartons Plant Breeders](#)



# Mendel and Darwin paved the way for scientific plant breeding




# The development of hybrid corn led to a big increase in yields




The progeny of two genetically different parents often show enhanced growth – this effect is termed “hybrid vigor”

Shull, G.H. (1909) A pure line method in corn breeding. Am. Breed. Assoc. Rep. 5, [51–59](#) by permission of Oxford University Press.

# Hybrid corn was rapidly adopted because of its increased yields



Even though farmers had to purchase seed every year, increased yields more than offset increased costs



Source: *Agricultural Statistics*, NASS, USDA, various years.

Shull, G.H. (1909) A pure line method in corn breeding. *Am. Breed. Assoc. Rep.* 5, 51–59 by permission of Oxford University Press; Economic Research Service / [USDA](#)



# Norman Borlaug was a plant breeder, and “father of the green revolution”



One of the most significant accomplishments of 20<sup>th</sup> century science was the development of lodging-resistant, high-yielding semi-dwarf grain varieties




Distinguished plant breeder and Nobel Laureate  
[Norman Borlaug](#) 1914-2009

# Improved green-revolution plants led to dramatically increased crop yields

The introduction of disease-resistant, semi-dwarf varieties turning countries from grain importers to grain exporters

Dwarf wheat was developed at **CIMMYT** – the International Maize and Wheat Improvement Center

Wheat yields in selected countries, 1950-2004



Source: FAO via [Brian0918](#)





Consultative Group on International Agricultural Research CGIAR

CGIAR is an international organization of agricultural research groups



IFPRI  
Wash. DC  
USA



International Center  
for Agricultural Research  
in the Dry Areas

ICARDA  
Aleppo  
Syrian Arab Rep.



ICRISAT  
Patancheru  
India

IRRI  
Los Baños  
Philippines



WorldFish  
Penang  
Malaysia



World Agroforestry Centre  
TRANSFORMING LIVES AND LANDSCAPES



CIP  
Lima  
Peru



AfricaRice



ILRI  
Nairobi  
Kenya

Africa Rice Center-WARDA  
Cotonou  
Benin

IITA  
Ibadan  
Nigeria

IWMI  
Colombo  
Sri Lanka

CIFOR  
Bogor  
Indonesia



Biodiversity  
International  
Rome  
Italy




AN INNOVATION FROM THE PLANT CELL

© 2011 American Society of Plant Biologists




# Rice breeding at IRRI also brought huge yield increases



IR8, released in 1966, “...was to tropical rices what the Model T Ford was to automobiles.” It was known as “miracle rice” because of its high yields.

Photo courtesy [IRRI](http://IRRI)

# Crop productivity has kept pace with population because of increased yields



**Growing more food without using more land helps mitigate climate change and slow the loss of biodiversity**

Burney, J.A., Davis, S.J., and Lobell, D.B. (2010). Greenhouse gas mitigation by agricultural intensification. *Proc. Natl. Acad. Sci.* 107: [12052-12057](https://doi.org/10.1073/pnas.1005210107).



# Modern plant breeders use molecular methods including DNA sequencing and proteomics as well as field studies



Photo credits Scott Bauer [USDA](#); [CIMMYT](#); [IRRI](#); [RCMI](#); [Duke](#) Institute for Genome Sciences and Policy




# Advances in genetic technologies contribute to improved plants

- Marker assisted selection
- Genome-wide association studies
- Recombinant DNA technology and transgenic plants




Photo credit: [IRRI](#)

# Marker assisted selection (MAS)




Phenotype: physical expression of traits



Genotype: sequence of all the genes in a genome


Photo credit [LemnaTec](#); Anderson, L.K., Lai, A., Stack, S.M., Rizzon, C. and Gaut, B.S. (2006). Uneven distribution of expressed sequence tag loci on maize pachytene chromosomes. *Genome Research*. 16: [115-122](#).

# Marker assisted selection (MAS)



Phenotype: physical expression of traits

Selecting for DNA markers is faster than selecting for phenotype




Genotype: sequence of all the genes in a genome

Photo credit [LemnaTec](#): Anderson, L.K., Lai, A., Stack, S.M., Rizzon, C. and Gaut, B.S. (2006). Uneven distribution of expressed sequence tag loci on maize pachytene chromosomes. *Genome Research*. 16: [115-122](#).



# How markers work: Each generation, genes reassort or shuffle



# Example: Introgression of a disease resistance gene




Elite tomato

We want to add a disease resistance trait to an “elite” tomato plant.



Poor tomato but **disease resistant**  
(resistance gene indicated)

# Example: Introgression of a disease resistance gene




We cross the two plants. Some of their progeny inherit the disease resistance trait, some don't – how can we tell the difference?

Photo by [Stephen Ausmus](#) USDA




# Example: Introgression of a disease resistance gene




We can use markers to look at their DNA and identify those with the resistance gene. It's faster and easier than infecting them to see the phenotype

# Example: Introgression of a disease resistance gene



Is this an elite, disease-resistant tomato? No, half of its genes are from the poor tomato


# Example: Introgression of a disease resistance gene



We have to repeatedly cross back to the elite tomato, using markers to identify plants with the disease resistance gene



# Example: Introgression of a disease resistance gene



Markers greatly accelerate breeding programs

After several generations, **elite, disease resistant** tomato

# MAS as a tool in production of submergence tolerant rice (*Sub1*)

Many rice-growing regions are prone to flooding. In Pakistan a 2010 a huge, deadly, flood submerged 17 million acres (69,000 km<sup>2</sup>) and destroyed much of the harvest






Photo credits: Abdul Majeed Goraya / [IRIN](#); [NASA](#) Goddard

# Submergence-tolerant rice can survive floods as long as 17 days


**Sensitive rice** – cannot survive prolonged flooding



FLOODING




Water retreats



**Submergence-tolerant *Sub1* rice** – growth arrests during flooding, enhancing survival




# Production of Swarna-Sub1: Cross Swarna with *Sub1* donor



Swarna –  
high  
yielding,  
flooding  
sensitive

Swarna-Sub1

Submergence  
tolerant parent



MAS allowed the *Sub-1* trait to be rapidly introgressed into Swarna. The Swarna-Sub1 rice accounted for over  $\frac{1}{4}$  of the rice planted in India in 2010.



Reprinted by permission from Macmillan Publishers Ltd. (NATURE) Xu, K., Xu, X., Fukao, T., Canlas, P., Maghirang-Rodriguez, R., Heuer, S., Ismail, A.M., Bailey-Serres, J., Ronald, P.C., and Mackill, D.J. (2006). Sub1A is an ethylene-response-factor-like gene that confers submergence tolerance to rice. Nature 442: [705-708](#). Photo courtesy of [Adam Barclay](#) CPS, IRRI Photo.







*Maize*



*Rice*



*Soy*




*Brassica*






**Genome sequence data are available for many important plants**



# Phenotype analysis



## Genotype analysis



## Gene discovery




## Association analysis



Genome-wide methods make it possible to identify genes associated with complex traits, like yield or water use efficiency

# This approach allows hundreds of genes with small effects to be identified


In maize, grain yields are correlated with leaf angle and size. A genome-wide association survey (GWAS) revealed hundreds of single-nucleotide polymorphisms (SNPs) associated with these traits, providing invaluable information for breeders.



Reprinted by permission from Macmillan Publishers Ltd. Tian, F., Bradbury, P.J., Brown, P.J., Hung, H., Sun, Q., Flint-Garcia, S., Rocheford, T.R., McMullen, M.D., Holland, J.B., and Buckler, E.S. (2011). Genome-wide association study of leaf architecture in the maize nested association mapping population. *Nat Genet* 43: [159-162](#).



# GWAS reveals SNPs that contribute to disease resistance



Similar studies have led to the identification of genes contributing to other agronomically important traits including drought tolerance

Reprinted by permission from Macmillan Publishers Ltd Kump, K.L., Bradbury, P.J., Wissler, R.J., Buckler, E.S., Belcher, A.R., Oropeza-Rosas, M.A., Zwonitzer, J.C., Kresovich, S., McMullen, M.D., Ware, D., Balint-Kurti, P.J., and Holland, J.B. (2011). Genome-wide association study of quantitative resistance to southern leaf blight in the maize nested association mapping population. *Nat Genet* 43: [163-168](#).

# Genetic Modification (GM) is another breeding method





Elite tomato

Recombinant DNA (or GM) allows a single gene to be introduced into a genome. This method can be faster than conventional breeding



Poor tomato but  
disease resistant

Elite, disease resistant tomato






# Why are GM methods used sometimes and molecular breeding others?

## Molecular breeding



1. Desired trait must be present in population



2. Genetic resources must be available




3. Plant should be propagated sexually

# Why are GM methods used sometimes and molecular breeding others?

## Molecular breeding



1. Desired trait must be present in population




2. Genetic resources must be available



3. Plant should be propagated sexually

## GM



# GM Example: Disease resistant banana by introduction of a gene from pepper



**Resistant**



**Susceptible**



# GM Example: Insect resistance through introduction of the *Bt* gene

Wild-type peanut plant



Peanut plant expressing the *Bt* gene



Photo by [Herb Pilcher](#) USDA









# GM Example: Herbicide resistance




Plants compete with other plants for sunlight and nutrients. Many farmers use herbicides to eliminate weeds (undesired plants) from their fields.

Left – corn rows sprayed with herbicide to eliminate competing plants  
Right – corn being choked by giant foxtail (*Setaria faberi*)

# Herbicide tolerant plants are environmentally friendly

Farmers that plant herbicide-tolerant crop plants use *less herbicide*, herbicides that are *less toxic*, and *till (plow) less*, saving soil and fuel.



Cerdeira, A.L. and Duke, S.O. (2006). The Current Status and Environmental Impacts of Glyphosate-Resistant Crops. *J. Environ. Qual.* 35: [1633-1658](#). Photo credit [Hunt Sanders](#), University of Georgia, [bugwood.org](#).



# Gene flow through pollen movement has to be monitored and controlled



There have been confirmed cases of gene transfer from crops to weeds and vice versa.

- What consequences are expected from gene flow?
- How can gene flow be minimized?
- How can consequences be mitigated?

# Future Challenges



Photo credit: [IRRI](#)



# Breeders can use more than one technology to address a challenge

Public – private partnerships

MAS breeding

Gene pyrimiding

SOLUTION

Improved agronomic practices

GM technology

Genome-wide association

PROBLEM



# Breeding plants for $\beta$ -carotene (pro-vitamin A) enrichment





Image sources: [Petaholmes](#) based on [WHO data](#);


# Enhanced $\beta$ -carotene content in food can prevent vitamin A deficiency

• Many staple foods are poor sources of  $\beta$ -carotene so many people do not get adequate vitamin A in their diet



# Synthesis, storage and breakdown all affect $\beta$ -carotene content

To increase beta-carotene levels in plants, you need **more synthesis**, **more storage** or **less catabolism**



**Chromoplasts** – organelles that store carotenoids





Photo credit: [University of Wisconsin](http://University of Wisconsin)





$\beta$ -carotene  
makes the rice  
look golden

# There is no inherently right or wrong way to enhance plant nutritional quality



$\beta$ -carotene



Vitamin A

The  $\beta$ -carotene  
enriched foods  
shown here  
have been  
produced using  
GM and non-GM  
approaches

Photo credit: [Golden rice humanitarian board](#)


# Biofortified plants are improving nutrition for many



The non-profit organization HarvestPlus focuses on the development of biofortified crops for the developing world, including a provitamin A enriched sweet potato that is currently being grown by **half a million** families. Other biofortification projects are underway to increase levels of protein, iron, zinc, antioxidants and other beneficial components in food.


Sources: [HarvestPlus](#); [CIMMYT](#)

# Breeding for drought tolerance






# Food production for one person for one day requires 3000 liters of water




[Comprehensive Assessment of Water Management in Agriculture](#). 2007. Water for Food, Water for Life: A Comprehensive Assessment of Water Management in Agriculture. London: Earthscan, and Colombo: International Water Management Institute.

# The incidence of major droughts is on the rise

China experienced a major drought in 2011



Major droughts and heat waves in China, Russia and Australia have impacted food production and raised prices



Russia experienced heat waves, drought and wildfires in 2010



# In 2011 seed companies released water-optimized corn

Both of these varieties were developed  
using modern molecular breeding methods  
without the use of recombinant DNA

 Agrisure Artesian















# Agricultural innovation in Africa – breeding crops for sub-Saharan Africa



Birth rates are high and crop yields are low so many African countries are not currently self-sufficient

# Many African countries experience a very high rate of undernourishment



Prevalence of undernourishment in developing countries (2005-07)

- Very high (undernourishment 35% and above)
- High (undernourishment 25-34%)
- Moderately high (undernourishment 15-24%)
- Moderately low (undernourishment 5-14%)
- Very low (undernourishment below 5%)
- Missing or insufficient data



www.fao.org

# The challenges to food production in Africa are immense

- Lack of infrastructure, especially irrigation and access to transportation networks
- High incidence of diseases
- Lack of available fertilizers
- Lack of education and support for farmers
- Lack of economic supports and market stability
- Agricultural subsidies in other countries affect market value



# Maize is a staple crop in Africa but very sensitive to drought damage

Less than 10% of crop land in sub-Saharan Africa is irrigated, making agriculture production highly susceptible to drought



Irrigation as percentage of cultivated area




Photo credit: Anne Wangalachi/[CIMMYT](#) Map Source – FAO [Aquastat](#) 2005

# As a consequence of climate changes, droughts are expected to increase

Drought events per country from 1970 to 2004 within Sub-Saharan Africa

In some African countries, yields from rain-fed agriculture, which is important for the poorest farmers, **could be reduced by up to 50% by 2020.**

-(FAO 2010)




Image credit: [United Nations Economic Commission for Africa](#), 2008 *Africa Review Report on Drought and Desertification*



# Water Efficient Maize for Africa was developed through a public-private partnership

Water-efficient maize optimized for growth in sub-Saharan Africa has been developed through a combination of breeding and GM methods



WEMA is being developed as a public-private partnership that includes international and regional plant breeding institutes, philanthropic groups and Monsanto

Photo credits: Anne Wangalachi/[CIMMYT](http://CIMMYT)



# Plant breeding can support African agriculture



African farmers need access to high yielding, drought tolerant, disease resistant plants. Most food is grown by small-scale farmers with little mechanization. Cassava, cowpea and banana are important crops and the focus of intensive breeding programs.



Photos courtesy of [IITA](#)

# African governments are working together to support agriculture

Alliance for a Green Revolution in Africa



*“AGRA is a dynamic, African-led partnership working across the African continent to help millions of small-scale farmers and their families lift themselves out of poverty and hunger”.* A major thrust of these efforts is to develop Africa’s human capacity through education, innovation and technology transfer.

Source: [AGRA](#)






**In the next 50 years, we will have to produce as much food as we have yet produced in human history**



# We have many paths to follow



# Breeding crops for a second green revolution



# Questions



# What risk assessments are performed on GM crops?

Before release into the environment, GM crops are subject to risk-assessment and risk-management measures to evaluate:

- Risks to human health (including toxicity and allergenicity)
- Risks of evolution of resistance in target pathogens or pests
- Risks to non-target organisms
- Risks from movement of transgenes



# Will genes from GMOs contaminate wild populations?

When Pandora opened the forbidden box she released evil into the world



John William Waterhouse: [Pandora](#) - 1896

# Will anti-insecticidal genes harm unintended targets?



ButterflyUtopia.com



*The evidence shows that the planting of GE crops has largely resulted in less adverse or equivalent effects on the farm environment compared with the conventional non-GE systems that GE crops replaced (National Academies 2010)*

Image credit [jons2](#)



# Will GMOs take away choice and exploit small farmers?

Insect Resistant  
Maize for Africa

> 45% of corn yields are often lost to insects

Partnerships including national agricultural research institutions, non-government and community-based organizations, regional research networks, and private companies work together to develop seeds that are suited to local conditions and are affordable for local farmers



Photo credit: [CIMMYT](#).

# Are GM crops safe to eat?

## YES

All GM plants are subject to extensive testing and regulatory oversight and no detrimental health effects have been identified



*Bt* corn is less prone to contamination by fungi which produce toxins linked to cancer and birth defects



GM biofortification can ensure that *all* children get adequate levels of protein, vitamins and mineral nutrients.



GM is a safe and beneficial tool in the quest to sustainably feed the growing population

# Scientists worldwide endorse GM as an important tool for breeding

*“Both genetic improvement and better crop management are vital and both should be resourced in parallel.” - 2009*



*“The ASPB believes strongly that, with continued responsible regulation and oversight, GE will bring many significant health and environmental benefits to the world and its people.” - 2006*

